

- 1739 *Epirrhoe rivata* Hb. Wood Carpet
Locally widespread in the southern half of England; larva feeds on Hedge and Lady's Bedstraw. Recorded in 1986.
- 1750 *Lampropteryx suffumata* D. & S. Water Carpet
Widely distributed and generally common throughout much of the British Isles; the species has been recorded in Devon during 1994; larva feeds on various species of Bedstraw. Recorded in 1983.
- 1757 *Eulithis mellinata* Fab. The Spinach
Widespread and moderately common in England and Wales; larva feeds on Red and Black Currant. Recorded in 1983.
- 1792 *Epirrita dilutata* D. & S. November Moth
Probably resident but until a full genitalia examination is carried out this will not be known; larva feeds on Blackthorn, Hawthorn and other species.
- 1807 *Perizoma albulata* D. & S. Grass Rivulet
Widespread and locally common; the foodplant, Yellow Rattle, is nearby. Recorded in 1989.
- 1827 *Eupithecia intricata* Zett. ssp. *arceuthata* Freyer. Freyer's Pug
13.6.1964 by A.H. Dobson & H.A. Kennard (*D.A. Ent. Section* 17th report: 47)
- 1844 *E. indigata* Hb. Ochreous Pug
Locally widespread but a possible vagrant from pinewoods nearby; larva feeds on Pine. Recorded in 1989.
- 1849 *E. fraxinata* Crewe. Ash Pug
The headquarters of this species are to be found in the south-east and east of the country where its larvae feed on Sea Buckthorn; there is an Ash feeding race which occurs in small numbers over the greater part of the British Isles; the species has been recorded in South Devon. Recorded in 1986.
- 1867 *Aplocera plagiata* Linn. Treble-bar
Found not uncommonly over the greater part of the British Isles, the species inhabits downland, moorland and woods; larva feeds on various species of St. Johns Wort. Found in card index.
- 1897 *Semiothisa wauaria* Linn. V-Moth
Widely distributed at low density; larva feeds on leaves of currant bushes. Recorded in 1983.
- 1905 *Pachynemina hippocastanaria* Hübn. Horse Chestnut
Well established and not uncommon in the south and south-west; a moth of heathland and it is resident on the local heathland; possibly a vagrant; larva feeds on heather. Found in card index.
- 1914 *Ennomos fuscantaria* Haw. Dusky Thorn
The species inhabits woodland and parkland and is not uncommon throughout England and Wales; can be confused with others of the same genus; larva feeds on Ash. Found in card index.
- 1926 *Apocheima pilosaria* D. & S. Pale Bridled Beauty
Generally distributed and common in England and Wales; the species has been recorded elsewhere; larva feeds on a variety of trees and shrubs. Recorded in 1985.
- 1932 *Agriopsis leucophaearia* D. & S. Spring Usher
Widespread and generally common over much of England and Wales; larva feeds on Oak and sometimes Apple. Recorded in 1984.
- 1934 *A. marginaria* Fab. Dotted Border
Widespread and more or less common over the greater part of the British Isles. The species has been seen in Teignmouth; larva feeds on a variety of trees and shrubs. Recorded in 1982.

- 1935 *Erannis defoliaria* Cl. Mottled Umber
The species is common almost everywhere; larva feeds on a variety of trees and shrubs. Recorded in 1985.
- 1943 *Boarmia roboraria* Thunb. Great Oak Beauty
A local species found not uncommonly in southern England including Devon; normally a species of Oak woodland; larva feeds on Oak. Recorded in 1984.
- 1944 *Serraca punctinalis* Scop. Pale Oak Beauty
A woodland species that is not uncommon in the south and east of England but is local in the south-west; worn specimens of this species could be confused with *P. rhomboidaria*, Willow Beauty; larva feeds on Oak, Birch and occasionally Sallow and Sycamore. Found in card index.

SPHINGIDAE

- 1986 *Hyles euphorbiae* Linn. Spurge Hawkmoth
"At Dawlish Warren on 1 July a male was found on palings and 5 July a fine female in a conservatory" (Lt. Col. H. Bridges) (French, 1949, Migration reports for 1948. *Entomologist* **82**: 106).

ARCTIIDAE

- 2049 *Eilema deplana* Esp. Buff Footman
An inhabitant of woodland and downland but the moth is local in southern and south-west England; easily confused with *E. griseola*, the Dingy Footman; larva feeds on lichens and algae growing on trees and shrubs. Found in card index.
- 2051 *Lithosia quadra* Linn. Four-spotted Footman
An uncommon but regular migrant which is also locally resident in mature woodland in south-west England where the larvae feed on lichens. Recorded in 1986.

NOCTUIDAE

- 2080 *Euxoa obelisca* D. & S. Square Spot Dart
Locally common around the coast from the Isle of Wight through Cornwall and into north Devon; possibility of confusion with the related *E. tritici*, the White-line Dart; the life history apparently unknown. Recorded in 1985.
- 2082 *E. nigricans* Linn. Garden Dart
Generally distributed throughout the British Isles but can be confused with *Xestia xanthographa*, the Square-spot Rustic. Larva feeds on a wide variety of wild and cultivated plants. Recorded in 1983.
- 2084 *Agrotis cinerea* D. & S. Light Feathered Rustic
Does occur in the south-west but is local and widespread; could have been misidentified with a banded form of *E. tritici*, the White-line Dart, or a dark variation of *A. clavis*, Heart & Club. Larva feeds on Wild Thyme and other low plants. Recorded in 1983.
- 2099 *Actebia praecox* Linn. Portland Moth
"Formerly taken at Dawlish Warren but although worked for has not been taken recently" (Stidson, 1951, *Lepidoptera of Devon*: 27); [1958] W.L. Coleridge (*D.A. Ent. Section* 11th report); larvae, 2.6.1957. G. Haggett (*D.A. Ent. Section*. 11th report).
- 2117 *Paradiarsia glareosa* Esp. Autumnal Rustic
Stitson, 1951, *Lepidoptera of Devon*: 27 Generally distributed throughout the British Isles; larva feeds on a wide variety of plants. Recorded in 1984.
- 2127 *Xestia ditrapezium* D. & S. Triple Spotted Clay
Locally widespread throughout much of Britain; can be confused with *X. triangulum*, the Double Square-spot larva feeds on Birch, Sallow, Bramble and other trees and shrubs. Recorded in 1986.
- 2130 *X. baja* D. & S. Dotted Clay
Generally distributed and often common throughout the British Isles where it inhabits heath and woodland; larva feeds on many trees and plants. Found in card index.

- 2133 *X. sexstrigata* Haw. Six-striped Rustic
Generally distributed throughout the British Isles; could be confused with *Diarsia rubi*, the Small Square-spot. Larva feeds on a variety of herbaceous plants. Recorded in 1986.
- 2135 *X. agathina* Dup. Heath Rustic
Widespread in the British Isles in suitable habitats; larva feeds on young shoots of heather. Recorded on 20.8.1987.
- 2136 *Naenia typica* Linn. The Gothic
Widespread over much of the British Isles but rarely common except in the Midlands; could be confused with *Tholera decimalis*, the Feathered Gothic. Larva feeds on a wide variety of trees and plants. Recorded in 1986.
- 2178 *Tholera decimalis* Poda. Feathered Gothic
Widespread and locally common over much of the British Isles; mutual confusion with *Naenia typica* is a possibility; larva feeds on a variety of grasses. Recorded in 1983.
- 2206 *Mythimna putrescens* Hb. Devonshire Wainscot
Very local in the south-west and rarely found outside Devon and Cornwall; usually a species of cliffs; larva feeds on coastal grasses. Recorded in 1984.
- 2214 *Cucullia chamomillae* D. & S. Chamomile Shark
E.D. Morgan and R.V. Solly (larvae), both in Stidson, *Lepidoptera of Devon*: 44; H.A. Kennard, 13.5.1961 (*D.A. Ent. Section* 14th report, p.117).
- 2208 *Mythimna loreyi* Dup. The Cosmopolitan
A migrant species which does appear along the south coast; larva feeds (in captivity) on Cock's-foot Grass. Recorded in 1984.
- 2247 *Dichonia aprilina* Linn. Merveille du Jour
Widespread and locally common in England and Wales in woodland and has been recorded at other sites in Devon; larva feeds on Oak. Recorded in 1982.
- 2250 *Blepharita adusta* Esp. Dark Brocade
Widely distributed, but local in the southern half of England but commoner in the north, the Midlands and Scotland. Probably not seen as the species can be confused with *brassicae* Cabbage Moth. A voucher would be needed. Recorded in 1983.
- 2252 *Polymixis flavicincta* D. & S. Large Ranunculus
Widespread and not uncommon in the southern half of England; recorded in Teignmouth and other sites; larva feeds on a wide variety of wild and cultivated plants. Recorded in 1985.
- 2260 *Conistra rubiginea* D. & S. Dotted Chestnut
A local woodland and wooded heathland species rarely found outside central southern and south-west England; larva feeds (in captivity) on leaves of fruit trees; has been found in the wild once on Apple. Recorded in 1986.
- 2277 *Moma alpium* Osb. Scarce Merveille du Jour
10.6.1956 by P. Crow per F.H. Lees (*D.A. Ent. Section* 9th report: 47).
- 2273 *Xanthia togata* Esp. Pink-barred Sallow
Widespread and moderately common over much of the British Isles; the species can be confused with *X. ictertia*, The Sallow; larva feeds on Sallow catkins and then low plants. Found in card index.
- 2300 *Mormo maura* Linn. Old Lady
Widespread and not uncommon in England; has been found in other localities during 1994. Larva feeds on a variety of trees and shrubs including Blackthorn, Hawthorn and Birch. Recorded in 1983.
- 2312 *Ipimorpha subtusa* D. & S. The Olive
Locally widespread throughout England and Wales but this specimen must have wandered; a voucher would be needed; larva feeds on Aspen and Poplar. Recorded in 1989.

- 2325 *Apamea oblonga* Haw. Crescent Striped
The V.C.H. give "taken at Dawlish Warren by Mr E.F. Studd" (Stidson, *Lepidoptera of Devon*: 33).
- 2331 *A. unanimitis* Hb. Small Clouded Brindle
Locally widespread in England, Wales and Scotland, inhabiting damp woodland, fenland and marshy places; this species is easily confused with *A. secalis*, the Common Rustic; larva feeds on various grasses. Found in card index.
- 2333 *A. anceps* D. & S. Large Nutmeg
Widespread and sometimes locally common in the southern half of England, except the south-west where it is local, less frequent and coastal; can be confused with large specimens of *Discestra trifolii*, The Nutmeg. Larva feeds on various grasses. Recorded in 1982.
- 2334 *A. sordens* Rustic Shoulder-knot
A widespread and usually common species throughout the British Isles but so far not recorded in 1994; larva feeds on Cock's-foot and other grasses. Recorded in 1986.
- 2335 *A. scolopacina* Esp. Slender Brindle
Locally widespread over much of England and Wales where it inhabits woodland; if the species was seen then it must have been a vagrant. This species can be confused with *A. secalis*; larva feeds on various grasses. Found in card index.
- 2358 *Amphipoea fucosa* Freyer Saltern Ear
A locally common and mainly coastal species sporadically distributed over much of Britain; larva feeds in the roots and bases of Annual Meadow-grass and other grasses. Recorded in 1982.
- 2360 *A. oculea* Linn. Ear Moth
Widespread and generally not uncommon over much of the British Isles; larva feeds in the roots and stems of Tufted-hair Grass and other grasses; and also stems of Butterbur. Recorded in 1982.
- 2371 *Archanara dissoluta* Treit. Brown-veined Wainscot
12.8.1961, H.A. Kennard (*D.A. Ent. Section* 14th report: 116).
- 2373 *A. sparganii* Esp. Webb's Wainscot
Well established in south-east England but more local in its range westward to Cornwall; quite possibly a vagrant; larva feeds in the stems of Common Reed-Mace. Found in card index.
- 2400 *Helicoverpa armigera* Hb. Scarce Bordered Straw
A migrant species which occurs from time to time along the south coast including Devon; is also imported with tomatoes and cut flowers; larva feeds on garden *Geranium* and other plants. Recorded in 1982.
- 2403 *Heliothis peltigera* D. & S. Bordered Straw
A migrant species which occurs more commonly than the previous species; larva breeds here and feeds on Sticky Groundsel, garden Marigolds and common Rest Harrow but cannot survive the winter. Recorded in 1986.
- 2410 *Lithacodia pygarga* Hufn. Marbled White-spot
Widespread and moderately common in the southern half of England and has been recorded in Devon during 1994; larva feeds on Purple Moor Grass and other grasses. Recorded in 1984.
- 2432 *Trichoplusia ni* Hb. The Ni Moth
A migrant species being an uncommon and irregular visitor, found mostly in southern England; larva feeds on Marigold and Sea Rocket, and in captivity Lettuce and Cabbage. Recorded in 1982.
- 2482 *Schrankia taenialis* Hb. White-line snout
Widely distributed, but local, over the southern half of England; the species has been seen at other sites, in Devon, during 1994; larval stages have not been found in the wild but in captivity will feed on Lettuce and flowers of heathers, and later on sliced runner beans. Recorded in 1984.

Part 3

A List of species which have been recorded from Dawlish Warren, but whose occurrence is thought unlikely.

In this brief listing, comments are made on species whose presence has been recorded either in the literature or in records kept by the Nature Reserve. All are felt to be erroneous and based upon misidentification.

- 1694 *Scopula ternata* Schrank. Smoky Wave
Exmoor is the best-known locality. Being a moorland species it is unlikely to have been found on Dawlish Warren. Possibly confused with *S. immutata*, Lesser-cream Wave. Recorded in 1983.
- 1701 *Idaea sylvestraria* Hb. Dotted-border Wave
In Devon, almost entirely restricted to well-established heathland. Possibly confused with *I. subsericeata*, Satin Wave. Recorded in 1983 and 1985.
- 1761 *Chloroclysta miata* Linn. Autumn-green Carpet
Recorded from Dartmoor. It is unlikely to be resident. Possibly confused with *C. siterata*, Red-green Carpet, which is known from the Warren. Recorded in 1982 and 1986.
- 1775 *Colostyia multistrigaria* Haw. Mottled Grey
Only one record made at the time that *T. carpinata*, Early Tooth-stripe was on the wing, and with which it was probably confused. Found in card index.
- 1815 *Eupithecia abietaria* Goeze. Cloaked Pug
A colony used to occur in the New Forest otherwise its headquarters is in Northumberland. Recorded in 1984.
- 1888 *Ligdia adustata* D. & S. Scorched Carpet
Probably confused with *C. cuculata*, Royal Mantle. Recorded in 1985.
- 1908 *Epione paralellaria* D. & S. Dark Bordered Beauty
A Yorkshire species and not known in the south. Probably misidentification of *E. repanaria*, Bordered Beauty. Recorded in 1984.
- 2031 *Leucoma salicis* Linn. White Satin
Likely to be a misidentification, or possibly a vagrant. Recorded in 1984.
- 2083 *Euxoa cursoria* Hufn. Coast Dart
Resident on the coast from Suffolk to Northumberland, and on the west coast from Cheshire to Cumberland. Possible confusion with either *E. tritici* White-line Dart or *Agrotis ripae* Sand Dart. Recorded in 1986 and 1989.
- 2104 *Standfussiana lucerneae* Hufn. Northern Rustic
A species of rocky cliffs which occurs from the Isle of Wight westwards to Cornwall and North Devon. Possibly a vagrant. Recorded in 1984.
- 2108 *Noctua orbona* Hufn. Lunar Yellow Underwing
The present strongholds of this species are in the Breck district of East Anglia, and the eastern border counties of England and Scotland. There are reports of this species occurring on Dartmoor and it is found on the Wiltshire Downs. Doubtless confused with *N. comes*. Recorded in 1984 and 1986.
- 2185 *Orthosia populeti* Fabr. Lead-coloured Drab
The foodplant, Aspen, is absent. Possibly confused with *O. incerta* Clouded Drab. Recorded in 1986.
- 2209 *Senta flammea* Curt. Flame Wainscot
This species occurs in the Cambridgeshire fenlands and Norfolk Broads; Probably confused with *Chilodes maritimus* ab. *wisemariensis* Silky Wainscot. Recorded in 1983.

- 2249 *Blepharita satura* D. & S. Beautiful Arches
Status of this species in Britain is poorly documented and confused by a plethora of doubtful records. Could be confused with *A. monoglypha* Dark Arches. Recorded in 1982.
- 2329 *Apamea furva britannica* Cock. The Confused
Very local in the south-west; in rocky places by the sea. This species is very easily confused with *A. remissa* Dusky Brocade. Recorded in 1983.
- 2332 *A. pabulatricula* Brahm. Union Rustic
Formerly resident but now considered extinct; used to be found in the midlands and north-east around Lincolnshire and Yorkshire. Possibly confused with a large specimen of *M. furuncula* Cloaked Minor. Recorded in 1983.
- 2346 *Photedes morrisii* Dale. Morris's Wainscot
Recorded in 1992 by Mr A. Kolaj but later found to be a worn *A. phragmitidis*.
- 2347 *P. extrema* Hb. Concolorous Wainscot
The main stronghold of this species is Huntingdonshire, Northamptonshire and mid-Lincolnshire. Possibly confused with *P. pygmina* Small Wainscot. Recorded in 1982 and other years.
- 2351 *P. brevilinea* Fenn. Fenn's Wainscot
This species is confined to East Anglia. Could be confused with *A. sparganii* Webb's Wainscot, but even this species is not that common in Devon. Recorded in 1984.
- 2494 *Paracolax derivalis* Hb. Clay Fan-foot
The species occurs in south-east England. Possibly confused with *C. liniarea* Clay-triple Lines. Recorded in 1984.

Index

| | | | | | |
|---------------------------|------|--------------------------|------|----------------------------|------|
| <i>abietaria</i> Goeze | 1815 | <i>areola</i> Esp. | 2243 | <i>caja</i> Linn. | 2057 |
| <i>abruptaria</i> Thunb. | 1936 | <i>argiolus</i> Linn. | 1580 | <i>cardamines</i> Linn. | 1553 |
| <i>absinthiata</i> Cl. | 1830 | <i>argus</i> Linn. | 1571 | <i>cardui</i> Linn. | 1591 |
| <i>advenella</i> Zinck. | 1439 | <i>armigera</i> Hb. | 2400 | <i>carpinata</i> Borkh. | 1881 |
| <i>adusta</i> Esp. | 2250 | <i>asinalis</i> Hb. | 1397 | <i>capucina</i> Linn. | 2008 |
| <i>adustata</i> D. & S. | 1888 | <i>atalanta</i> Linn. | 1590 | <i>centaureata</i> D. & S. | 1825 |
| <i>aegaria</i> Linn. | 1614 | <i>augur</i> Fabr. | 2114 | <i>cerasi</i> Fabr. | 2187 |
| <i>aescularia</i> D. & S. | 1663 | <i>aurantiaria</i> Hb. | 1933 | <i>cespitis</i> D. & S. | 2177 |
| <i>aestivaria</i> Hb. | 1669 | <i>aversata</i> Linn. | 1713 | <i>chanomillae</i> D. & S. | 2214 |
| <i>agathina</i> Dup. | 2135 | <i>badiata</i> D. & S. | 1746 | <i>chenopodiata</i> Linn. | 1732 |
| <i>agestis</i> D. & S. | 1572 | <i>baja</i> D. & S. | 2130 | <i>chlorosata</i> Scop. | 1902 |
| <i>albicolon</i> Hb. | 2152 | <i>batis</i> Linn. | 1652 | <i>christyi</i> Allen | 1796 |
| <i>albula</i> D. & S. | 2076 | <i>bembiciformis</i> Hb. | 0371 | <i>chrysitis</i> Linn. | 2434 |
| <i>albulata</i> D. & S. | 1807 | <i>betularia</i> Linn. | 1931 | <i>chrysoprasaria</i> Esp. | 1673 |
| <i>alchemillata</i> Linn. | 1803 | <i>bicolorata</i> Hufn. | 2164 | <i>chrysorrhoea</i> Linn. | 2029 |
| <i>alni</i> Linn. | 2281 | <i>bicruris</i> Hufn. | 2173 | <i>cinerea</i> D. & S. | 2084 |
| <i>alniaria</i> Linn. | 1913 | <i>bidentata</i> Cl. | 1920 | <i>circellaris</i> Hufn. | 2262 |
| <i>alpinella</i> Hb. | 1325 | <i>bilineata</i> Linn. | 1742 | <i>citrata</i> Linn. | 1762 |
| <i>alpium</i> Osb. | 2227 | <i>binaria</i> Hufn. | 1646 | <i>clavipalpis</i> Scop. | 2389 |
| <i>alsines</i> Brahm. | 2381 | <i>biselata</i> Hufn. | 1702 | <i>clavis</i> Hufn. | 2088 |
| <i>alternata</i> Mull. | 1738 | <i>bistortata</i> Goeze | 1947 | <i>clorana</i> Linn. | 2418 |
| <i>alternaria</i> Hb. | 1890 | <i>blanda</i> D. & S. | 2382 | <i>c-nigrum</i> Linn. | 2126 |
| <i>ambigua</i> D. & S. | 2384 | <i>brassicae</i> Linn. | 1549 | <i>comes</i> Hb. | 2109 |
| <i>anceps</i> D. & S. | 2333 | <i>brassicae</i> Linn. | 2154 | <i>comma</i> Hb. | 2205 |
| <i>anceps</i> Goeze | 2005 | <i>brevilinea</i> Fenn | 2351 | <i>complana</i> Linn. | 2047 |
| <i>angustea</i> Curt. | 1342 | <i>britannica</i> Turn. | 1769 | <i>confusa</i> Hufn. | 2171 |
| <i>antiopa</i> Linn. | 1596 | <i>brumata</i> Linn. | 1799 | <i>conigera</i> D. & S. | 2192 |
| <i>antiqua</i> Linn. | 2026 | <i>bucephala</i> Linn. | 1994 | <i>contaminella</i> Hb. | 1323 |
| <i>aprilina</i> Linn. | 2247 | <i>c-album</i> L. | 1598 | <i>convolvuli</i> Linn. | 1972 |

| | | | | | |
|------------------------------|------|------------------------------|------|------------------------------|------|
| <i>coronata</i> Hufn. | 1378 | <i>forficalis</i> Linn. | 1356 | <i>lichenaria</i> Hufn. | 1945 |
| <i>costalis</i> Fabr. | 1413 | <i>forficella</i> Thunb. | 1329 | <i>lichenea</i> Hb. | 2255 |
| <i>coryli</i> Linn. | 2425 | <i>fraxinata</i> Crewe | 1849 | <i>ligustri</i> D. & S. | 2291 |
| <i>crenata</i> Hufn. | 2326 | <i>fucosa</i> Freyer | 2358 | <i>linariata</i> D. & S. | 1816 |
| <i>crepuscularia</i> D. & S. | 1948 | <i>fuliginosa</i> Linn. | 2064 | <i>literosa</i> Haw. | 2342 |
| <i>cribrella</i> Hb. | 1458 | <i>fulvata</i> Forst. | 1765 | <i>lithoxylea</i> D. & S. | 2322 |
| <i>crocealis</i> Hb. | 1385 | <i>furcata</i> Thunb. | 1777 | <i>litoralis</i> Curt. | 2201 |
| <i>croceus</i> Geoffr. | 1545 | <i>furcula</i> Cl. | 1997 | <i>liturata</i> Cl. | 1893 |
| <i>cruda</i> D. & S. | 2182 | <i>furuncula</i> D. & S. | 2341 | <i>loreyi</i> Dup. | 2208 |
| <i>cuculata</i> Hufn. | 1736 | <i>furva</i> D. & S. | 2329 | <i>lota</i> Cl. | 2263 |
| <i>cucullatella</i> Linn. | 2077 | <i>fuscantaria</i> Haw. | 1914 | <i>lotella</i> Hb. | 1432 |
| <i>culmella</i> Linn. | 1293 | <i>fuscovenosa</i> Goeze | 1705 | <i>lubricipeda</i> Linn. | 2060 |
| <i>cultraria</i> Fabr. | 1647 | <i>galathea</i> Linn. | 1620 | <i>lucerna</i> Linn. | 2104 |
| <i>cursoria</i> Hufn. | 2083 | <i>galiata</i> D. & S. | 1740 | <i>lucipara</i> Linn. | 2305 |
| <i>curtula</i> Linn. | 2019 | <i>gamma</i> Linn. | 2441 | <i>lunaedactyla</i> Linn. | 1495 |
| <i>decimalis</i> Poda | 2748 | <i>geminipuncta</i> Haw. | 2370 | <i>lunosa</i> Haw. | 2270 |
| <i>defoliaria</i> Cl. | 1935 | <i>geniculea</i> Haw. | 1309 | <i>lupulinus</i> Linn. | 0017 |
| <i>dentaria</i> Fabr. | 1917 | <i>genistella</i> Dup. | 1443 | <i>luridata</i> Hufn. | 1734 |
| <i>deplana</i> Esp. | 2049 | <i>glareosa</i> Esp. | 2117 | <i>lurideola</i> Zinck. | 2050 |
| <i>derivialis</i> Hb. | 2494 | <i>gnoma</i> Fabr. | 2006 | <i>luteolata</i> Linn. | 1906 |
| <i>derivata</i> D. & S. | 1747 | <i>gothica</i> Linn. | 2190 | <i>lutea</i> Hufn. | 2061 |
| <i>designata</i> Hufn. | 1722 | <i>gracilis</i> D. & S. | 2186 | <i>lutosa</i> Hubn. | 2375 |
| <i>didyma</i> Esp. | 2343 | <i>graminis</i> Linn. | 2176 | <i>lychnidis</i> D. & S. | 2267 |
| <i>dilatata</i> D. & S. | 1795 | <i>griseata</i> Peters | 1682 | <i>macilenta</i> Hb. | 2264 |
| <i>dimidiata</i> Hufn. | 1708 | <i>grisealis</i> D. & S. | 2492 | <i>marginepunctata</i> Goeze | 1689 |
| <i>dissoluta</i> Treit. | 2371 | <i>griseola</i> Hb. | 2044 | <i>margariata</i> Linn. | 1961 |
| <i>ditrapezium</i> D. & S. | 2127 | <i>grossulariata</i> Linn. | 1884 | <i>marginaria</i> Fabr. | 1934 |
| <i>dodonaea</i> D. & S. | 2014 | <i>halterata</i> Hufn. | 1879 | <i>marginaria</i> Linn. | 1887 |
| <i>dominula</i> Linn. | 2068 | <i>haworthiata</i> Doubl. | 1813 | <i>maritimus</i> Tausch. | 2391 |
| <i>dromedarius</i> Linn. | 2000 | <i>hippocastanaria</i> Hubn. | 1905 | <i>matura</i> Hufn. | 2303 |
| <i>duplaris</i> Linn. | 1657 | <i>hortulata</i> Linn. | 1376 | <i>maura</i> Linn. | 2300 |
| <i>elinguarua</i> Linn. | 1921 | <i>humuli</i> Linn. | 0014 | <i>megacephala</i> D. & S. | 1278 |
| <i>elpenor</i> Linn. | 1991 | <i>icarus</i> Rott. | 1574 | <i>megea</i> Linn. | 1615 |
| <i>emarginata</i> Linn. | 1712 | <i>icterata</i> Vill. | 1838 | <i>mellinata</i> Fabr. | 1757 |
| <i>epomidion</i> Haw. | 2327 | <i>icteritia</i> Hufn. | 2274 | <i>mendica</i> Fabr. | 2120 |
| <i>eremita</i> Fabr. | 2248 | <i>imitaria</i> Hb. | 1690 | <i>meticulosa</i> Linn. | 2306 |
| <i>erosaria</i> D. & S. | 1915 | <i>immutata</i> Linn. | 1692 | <i>mercurella</i> Linn. | 1344 |
| <i>euphoibiae</i> L. | 1986 | <i>impluviata</i> D. & S. | 1778 | <i>miata</i> Linn. | 1761 |
| <i>exanthemata</i> Scop. | 1956 | <i>impura</i> Hb. | 2198 | <i>micacea</i> Esp. | 2361 |
| <i>exclamationis</i> Linn. | 2089 | <i>incerta</i> Hufn. | 2188 | <i>miniata</i> Forst. | 2037 |
| <i>exigua</i> Hb. | 2385 | <i>indigata</i> Hb. | 1844 | <i>minima</i> Haw. | 2345 |
| <i>expallidata</i> Doubl. | 1833 | <i>inquinatella</i> D. & S. | 1306 | <i>multistrigaria</i> Haw. | 1775 |
| <i>extrema</i> Hb. | 2347 | <i>interjecta</i> Hb. | 2112 | <i>monacha</i> Linn. | 2033 |
| <i>falcataria</i> Linn. | 1648 | <i>intricata</i> Zell. | 1827 | <i>monodactyla</i> Linn. | 1524 |
| <i>fasciaria</i> Linn. | 1962 | <i>io</i> Linn. | 1597 | <i>monoglypha</i> Hufn. | 2321 |
| <i>fasciuncula</i> Haw. | 2340 | <i>ipsilon</i> Hufn. | 2091 | <i>montanata</i> D. & S. | 1727 |
| <i>ferrago</i> Fabr. | 2193 | <i>jacobaeae</i> Linn. | 2069 | <i>morpheus</i> Hufn. | 2387 |
| <i>festucae</i> Linn. | 2439 | <i>janthe</i> D. & S. | 2111 | <i>morrisii</i> Dale | 2346 |
| <i>ferrugata</i> Cl. | 1725 | <i>jota</i> Linn. | 2443 | <i>muralis</i> Forst. | 2295 |
| <i>ferrugalis</i> Hb. | 1395 | <i>jurtina</i> Linn. | 1626 | <i>napi</i> Linn. | 1551 |
| <i>ferruginea</i> Esp. | 2302 | <i>lacertinaria</i> Linn. | 1645 | <i>nana</i> Hufn. | 2147 |
| <i>filipendulae</i> Linn. | 0169 | <i>lactearia</i> Linn. | 1674 | <i>nanata</i> Hb. | 1846 |
| <i>fimbriata</i> Schreb. | 2110 | <i>lacustrata</i> Panzer | 1338 | <i>nebulata</i> Scop. | 1874 |
| <i>flammea</i> Curt. | 2209 | <i>l-album</i> Linn. | 2202 | <i>nebulosa</i> Hufn. | 2150 |
| <i>flammealis</i> D. & S. | 1424 | <i>latistria</i> Haw. | 1307 | <i>neustria</i> Linn. | 1634 |
| <i>flavago</i> D. & S. | 2364 | <i>latruncula</i> D. & S. | 2339 | <i>ni</i> Hb. | 2432 |
| <i>flavicincta</i> D. & S. | 2252 | <i>legatella</i> D. & S. | 1866 | <i>nigra</i> Haw. | 2232 |
| <i>flavofasciata</i> Thunb. | 1808 | <i>leporina</i> Linn. | 2280 | <i>nigricans</i> Linn. | 2082 |
| <i>flexula</i> D. & S. | 2473 | <i>leucophaearia</i> D. & S. | 1932 | <i>noctuella</i> D. & S. | 1398 |
| <i>fluctuata</i> Linn. | 1728 | <i>libatrix</i> Linn. | 2469 | <i>nubilalis</i> Hb. | 1375 |

| | | | | | |
|--|------|---|------|-----------------------------|------|
| <i>nupta</i> Linn. | 2452 | <i>puta</i> Hb. | 2092 | <i>straminella</i> D. & S. | 1304 |
| <i>obelisca</i> D. & S. | 2080 | <i>putrescens</i> Hb. | 2206 | <i>strigilis</i> Linn. | 2337 |
| <i>obeliscata</i> Hb. | 1768 | <i>putris</i> Linn. | 2098 | <i>suasa</i> D. & S. | 2159 |
| <i>oblonga</i> Haw. | 2325 | <i>pygarga</i> Hufn. | 2410 | <i>subfusca</i> Haw. | 1332 |
| <i>obstipata</i> Fabr. | 1719 | <i>pygmina</i> Haw. | 2350 | <i>subfuscata</i> Haw. | 1837 |
| <i>ocellata</i> Linn. | | <i>pyraliata</i> D. & S. | 1758 | <i>subsericeata</i> Haw. | 1709 |
| (<i>Cosmorhoe</i>) | 1752 | <i>pyrina</i> Linn. | 0161 | <i>subtusa</i> D. & S. | 2312 |
| <i>ocellata</i> Linn. | | <i>pyritoides</i> Hufn. | 1653 | <i>succenturiata</i> Linn. | 1839 |
| (<i>Smerinthus</i>) | 1980 | <i>quadra</i> Linn. | 2051 | <i>suffumata</i> D. & S. | 1750 |
| <i>ochrearia</i> Rossi. | 1968 | <i>quadrifasiata</i> Cl. | 1726 | <i>sylvestriaria</i> Hb. | 1701 |
| <i>ochroleuca</i> D. & S. | 2352 | <i>quadripunctaria</i> Poda | 2067 | <i>sylvestris</i> Poda | 1526 |
| <i>ocularis</i> Linn. | 1654 | <i>quercinaria</i> Hufn. | 1912 | <i>syringaria</i> Linn. | 1910 |
| <i>oculea</i> Linn. | 2360 | <i>quercus</i> Linn. | | <i>taenialis</i> Hb. | 2482 |
| <i>odites</i> Hb. | 2226 | (<i>Lasiocampa</i>) | 1637 | <i>tarsipennalis</i> Treit. | 2489 |
| <i>oleracea</i> Linn. | 2160 | <i>quercus</i> Linn. (<i>Quercusia</i>) | 1557 | <i>temerata</i> D. & S. | 1958 |
| <i>olivata</i> D. & S. | 1774 | <i>rapae</i> Linn. | 1550 | <i>tenuiata</i> Hb. | 1811 |
| <i>ophiogramma</i> Esp. | 2336 | <i>rectangulata</i> Linn. | 1860 | <i>ternata</i> Schrank | 1694 |
| <i>orbona</i> Hufn. | 2108 | <i>remissa</i> Hb. | 2330 | <i>testacea</i> D. & S. | 2353 |
| <i>oxycanthea</i> Linn. | 2245 | <i>repandaria</i> Hufn. | 1907 | <i>tetralunaria</i> Hufn. | 1919 |
| <i>pabulatricula</i> Brahm | 2332 | <i>repandata</i> Linn. | 1941 | <i>thalassina</i> Hufn. | 2158 |
| <i>pallens</i> Linn. | 2199 | <i>rhamni</i> Linn. | 1546 | <i>tithonus</i> Linn. | 1625 |
| <i>pallida</i> Curt. | 1336 | <i>rhomboidaria</i> D. & S. | 1937 | <i>togata</i> Esp. | 2273 |
| <i>palpina</i> Cl. | 2011 | <i>ribcata</i> Cl. | 1940 | <i>tragopoginis</i> Cl. | 2299 |
| <i>papilionaria</i> Linn. | 1666 | <i>ripae</i> Hb. | 2093 | <i>transversata</i> Hufn. | 1792 |
| <i>paralellaria</i> D. & S. | 1908 | <i>rivata</i> Hb. | 1739 | <i>trapezina</i> Linn. | 2318 |
| <i>pascuella</i> Linn. | 1294 | <i>rivularis</i> Fabr. | 2166 | <i>tremula</i> Cl. | 2007 |
| <i>pectinataria</i> Knoch | 1776 | <i>roboraria</i> Thunb. | 1943 | <i>triangulum</i> Hufn. | 2128 |
| <i>peltigera</i> D. & S. | 2403 | <i>roborella</i> D. & S. | 1452 | <i>tridens</i> D. & S. | 2283 |
| <i>pennaria</i> Linn. | 1923 | <i>ruberata</i> Freyer | 1779 | <i>trifolii</i> Hufn. | 2145 |
| <i>perlella</i> Scop. | 1302 | <i>rubi</i> Linn. | 1555 | <i>trifolii</i> Esp. | 0170 |
| <i>persicariae</i> Linn. | 2155 | <i>rubi</i> View. | 2123 | <i>trigrammica</i> Hufn. | 2380 |
| <i>phlaea</i> Linn. | 1561 | <i>rubidata</i> D. & S. | 1735 | <i>triplasia</i> Linn. | 2450 |
| <i>phoeniciata</i> Ramb. | 1855 | <i>rubiginea</i> D. & S. | 2260 | <i>tristella</i> D. & S. | 1305 |
| <i>phragmitidis</i> Hb. | 2377 | <i>rufa</i> Haw. | 2379 | <i>tritici</i> Linn. | 2081 |
| <i>pilosaria</i> D. & S. | 1926 | <i>ruffifasciata</i> Haw. | 1862 | <i>truncata</i> Hufn. | 1764 |
| <i>pinella</i> Linn. | 1313 | <i>rumicis</i> Linn. | 2289 | <i>trux</i> Hb. | 2090 |
| <i>psi</i> Linn. | 2163 | <i>ruralis</i> Scop. | 1405 | <i>turca</i> L. | 2191 |
| <i>plagiata</i> Linn. | 1867 | <i>sacraria</i> Linn. | 1716 | <i>typica</i> Linn. | 2136 |
| <i>plecta</i> Linn. | 2102 | <i>salicis</i> Linn. | 2031 | <i>umbra</i> Hufn. | 2399 |
| <i>porata</i> Linn. | 1679 | <i>sambucaria</i> Linn. | 1922 | <i>umbratica</i> Linn. | 2216 |
| <i>populeti</i> Fabr. | 2185 | <i>satura</i> D. & S. | 2249 | <i>unanimis</i> Hb. | 2331 |
| <i>populi</i> Linn. (<i>Laothoe</i>) | 1981 | <i>saucia</i> Hb. | 2119 | <i>undulata</i> Linn. | 1789 |
| <i>populi</i> Linn. | | <i>saxicola</i> Vaugh. | 1484 | <i>urticae</i> Linn. | 1593 |
| (<i>Poecilocampa</i>) | 1631 | <i>scolopacina</i> Esp. | 2335 | <i>vaccinii</i> Linn. | 2258 |
| <i>porcellus</i> Linn. | 1992 | <i>secalis</i> Linn. | 2343 | <i>v-ata</i> Haw. | 1858 |
| <i>porphyrea</i> D. & S. | 2118 | <i>segetum</i> D. & S. | 2087 | <i>venata</i> Brem. & Grey | 1531 |
| <i>potatoria</i> Linn. | 1640 | <i>semele</i> D. & S. | 1621 | <i>vestigialis</i> Hufn. | 2085 |
| <i>praecox</i> L. | 2099 | <i>sericealis</i> Scop. | 2474 | <i>versicolor</i> Borkh. | 2338 |
| <i>prasina</i> D. & S. | 2138 | <i>sexalata</i> Retz. | 1882 | <i>villica</i> Linn. | 2058 |
| <i>proboscidalis</i> Linn. | 2477 | <i>sexstrigata</i> Haw. | 2133 | <i>viminalis</i> Fabr. | 2225 |
| <i>pronuba</i> Linn. | 2107 | <i>silaceata</i> D. & S. | 1759 | <i>viretata</i> Hb. | 1883 |
| <i>pruinata</i> Hufn. | 1665 | <i>similis</i> Fuess. | 2030 | <i>virgata</i> Hufn. | 1718 |
| <i>prunalis</i> D. & S. | 1390 | <i>siterata</i> Hufn. | 1760 | <i>viridaria</i> Cl. | 2470 |
| <i>prunata</i> Linn. | 1754 | <i>sociella</i> Linn. | 1428 | <i>vitalbata</i> D. & S. | 1781 |
| <i>psi</i> Linn. | 2284 | <i>sordens</i> Hufn. | 2334 | <i>vulgata</i> Haw. | 1834 |
| <i>pudibunda</i> Linn. | 2028 | <i>spadicearia</i> D. & S. | 1724 | <i>w-album</i> Knoch | 1558 |
| <i>punctalis</i> Fabr. | 1414 | <i>sparganii</i> Esp. | 2373 | <i>wauria</i> Linn. | 1897 |
| <i>punctaria</i> Linn. | 1680 | <i>stagnata</i> Don. | 1350 | <i>w-latinum</i> Hufn. | 2157 |
| <i>punctinalis</i> Scop. | 1944 | <i>stellatarum</i> Linn. | 1984 | <i>xanthographa</i> D. & S. | 2134 |
| <i>pusaria</i> Linn. | 1955 | <i>straminea</i> Treit. | 2197 | <i>ziczac</i> Linn. | 2003 |